



TITLE:

THE FIXED POINT PROPERTY OF \$A\$-DIRECT SUMS OF \$N\$ UNIFORMLY NON-SQUARE BANACH SPACES (Nonlinear Analysis and Convex Analysis)

AUTHOR(S):

Tamura, Takayuki; Kato, Mikio

CITATION:

Tamura, Takayuki ...[et al]. THE FIXED POINT PROPERTY OF \$A\$-DIRECT SUMS OF \$N\$ UNIFORMLY NON-SQUARE BANACH SPACES (Nonlinear Analysis and Convex Analysis). 数理解析研究所講究録 2019, 2114: 133-136

ISSUE DATE:

2019-05

URL:

<http://hdl.handle.net/2433/252046>

RIGHT:

THE FIXED POINT PROPERTY OF A-DIRECT SUMS OF N UNIFORMLY NON-SQUARE BANACH SPACES

TAKAYUKI TAMURA

Graduate School of Social Sciences, Chiba University
ttakayuki@faculty.chiba-u.jp and

MIKIO KATO

Kyushu Institute of Technology
katom@mns.kyutech.ac.jp

Abstract

We shall show the fixed point property of A -direct sums of N uniformly non-square Banach spaces by characterizing the nontrivialness of Dominguez-Benavides coefficient $R(1, (X_1 \oplus \cdots \oplus X_N)_A)$, that is, $R(1, (X_1 \oplus \cdots \oplus X_N)_A) < 2$.

A norm $\|\cdot\|$ on \mathbb{R}^N is called *monotone* if $\|\mathbf{a}\| \leq \|\mathbf{b}\|$ for all $\mathbf{a} = (a_j), \mathbf{b} = (b_j) \in \mathbb{R}^N$ with $|a_j| \leq |b_j|$ ($j = 1, \dots, N$). For $\mathbf{a} = (a_j)$ $|\mathbf{a}|$ is defined by $|\mathbf{a}| = (|a_j|) \in \mathbb{R}^N$. A norm $\|\cdot\|$ on \mathbb{R}^N is called *absolute* if $\|\mathbf{a}\| = \| |\mathbf{a}| \|$ for all $\mathbf{a} \in \mathbb{R}^N$ and *normalized* if $\|e_j\| = 1$ for all $1 \leq j \leq N$, where e_j is the j -th unit vector in \mathbb{R}^N .

In [4] and [10] A -direct sums and AN -direct sums of N Banach spaces were introduced respectively by the following: Let $\|\cdot\|_A$ be an *arbitrary* norm on \mathbb{R}^N . The *A-direct sum* $(X_1 \oplus \cdots \oplus X_N)_A$ is the direct sum of X_1, \dots, X_N equipped with the norm

$$\|(x_1, \dots, x_N)\|_A = \|(\|x_1\|, \dots, \|x_N\|)\|_A, (x_1, \dots, x_N) \in X_1 \oplus \cdots \oplus X_N$$

and an *AN-direct sum* is an A -direct sum whose norm is defined from some absolute normalized norm $\|\cdot\|_{AN}$ on \mathbb{R}^N . It is known that a norm $\|\cdot\|_A$ on \mathbb{R}^N is absolute if and only if it is monotone ([2],[4],[14]).

In [13] Z -direct sums were introduced by the following: Let $\|\cdot\|_Z$ be an *monotone* norm on \mathbb{R}_+^N . The *Z-direct sum* $(X_1 \oplus \cdots \oplus X_N)_Z$ is the direct sum of X_1, \dots, X_N equipped with the norm

$$\|(x_1, \dots, x_N)\|_Z = \|(\|x_1\|, \dots, \|x_N\|)\|_Z, (x_1, \dots, x_N) \in X_1 \oplus \cdots \oplus X_N$$

Then we see that Z -direct sum and AN -direct sum are A -direct sum. Since an A -direct sum is isometric isomorphic to some AN -direct sum ([4]), then we have the following theorem.

Theorem 1 (cf. [4]). *Let X_1, \dots, X_N be Banach spaces. Let $\|\cdot\|_A$ be an arbitrary norm on \mathbb{R}^N . Then the norm of $(X_1 \oplus \cdots \oplus X_N)_A$ is monotone, that is,*

$$\|(x_1, \dots, x_N)\|_A \leq \|(y_1, \dots, y_N)\|_A$$

for $(x_1, \dots, x_N), (y_1, \dots, y_N) \in (X_1 \oplus \dots \oplus X_N)_A$ with $\|x_j\| \leq \|y_j\|$ ($j = 1, \dots, N$).

As usual S_X and B_X stand for the unit sphere and the closed unit ball of X , respectively. A Banach space X is said to have the *fixed point property* (resp. *weak fixed point property*) for nonexpansive mappings if every nonexpansive self-mapping T of any nonempty bounded closed (resp. weakly compact) convex subset C of X has a fixed point (T is called *nonexpansive* if $\|Tx - Ty\| \leq \|x - y\|$ for all $x, y \in C$). In [6] the coefficient $R(a, X)$ called as Dominguez-Benavides coefficient (cf. [3]) was introduced by the following: For $0 \leq a \leq 1$ let

$$R(a, X) = \sup \left\{ \liminf_{n \rightarrow \infty} \|x_n + x\| \right\},$$

where the supremum is taken over all $x \in X$ with $\|x\| \leq a$ and all weakly null sequences $\{x_n\}_n$ in the unit ball of X such that

$$\lim_{n, m \rightarrow \infty; n \neq m} \|x_n - x_m\| \leq 1.$$

In this paper we shall show the fixed point property for nonexpansive mappings of A -direct sums of N uniformly non-square Banach spaces by characterizing the nontrivialness of Dominguez-Benavides coefficient $R(1, (X_1 \oplus \dots \oplus X_N)_A)$, that is, $R(1, (X_1 \oplus \dots \oplus X_N)_A) < 2$.

The following theorem was proved in [6].

Theorem 2 ([6]). *Let X be a Banach space. If $R(a, X) < 1 + a$ for some $a > 0$, then X has the weak fixed point property for nonexpansive mappings.*

A Banach space X is called *uniformly non-square* ([9]) if there exists a constant $\varepsilon > 0$ such that

$$\min\{\|x + y\|, \|x - y\|\} \leq 2(1 - \varepsilon) \quad \text{for all } x, y \in S_X.$$

By Theorem 2 García-Falset et. al. [8] obtained the following remarkable result.

Theorem 3 ([8]). *Let X be a uniformly non-square Banach space. Then $R(1, X) < 2$ and hence X has the fixed point property for nonexpansive mappings.*

In [7] the following notions were introduced.

Definition 4 ([7]). For $\mathbf{a} = (a_j) \in \mathbb{R}^N$ let $\text{supp } \mathbf{a} = \{j : a_j \neq 0\}$.

(i) A norm $\|\cdot\|$ on \mathbb{R}^N is said to have *Property T_1^N* if

$$\|\mathbf{a}\| = \|\mathbf{b}\| = \frac{1}{2}\|\mathbf{a} + \mathbf{b}\| = 1, \quad \mathbf{a}, \mathbf{b} \in \mathbb{R}^N \implies \text{supp } \mathbf{a} \cap \text{supp } \mathbf{b} \neq \emptyset.$$

(ii) A norm $\|\cdot\|$ on \mathbb{R}^N is said to have *Property T_∞^N* if

$$\|\mathbf{a}\| = \|\mathbf{b}\| = \|\mathbf{a} + \mathbf{b}\| = 1 \implies \text{supp } \mathbf{a} \cap \text{supp } \mathbf{b} \neq \emptyset.$$

To show our key result we need the following propositions.

Proposition 5 ([11]). *Let $\{x_n^{(k)}\}_{n,k}, \{y_n^{(k)}\}_{n,k}$ be double sequences with nonzero terms in a Banach space X such that*

$$\lim_{k \rightarrow \infty} \liminf_{n \rightarrow \infty} \|x_n^{(k)}\| > 0, \lim_{k \rightarrow \infty} \liminf_{n \rightarrow \infty} \|y_n^{(k)}\| > 0.$$

Then the following are equivalent.

$$(i) \lim_{k \rightarrow \infty} \liminf_{n \rightarrow \infty} \|x_n^{(k)} + y_n^{(k)}\| = \lim_{k \rightarrow \infty} \liminf_{n \rightarrow \infty} (\|x_n^{(k)}\| + \|y_n^{(k)}\|).$$

$$(ii) \lim_{k \rightarrow \infty} \liminf_{n \rightarrow \infty} \left\| \frac{x_n^{(k)}}{\|x_n^{(k)}\|} + \frac{y_n^{(k)}}{\|y_n^{(k)}\|} \right\| = 2.$$

Proposition 6 ([5]; see also [1, Chpter III, Theorem 1.5]). *Let $\{x_n\}$ be a bounded sequence in a Banach space X . Then $\{x_n\}$ contains a subsequence $\{x_{n_k}\}$ such that $\lim_{k,l \rightarrow \infty; k \neq l} \|x_{n_k} - x_{n_l}\|$ exists.*

Proposition 7 ([16]). *Let $\{x_n\}$ be a weakly null sequence in a Banach space X . Assume that $\lim_{n,m \rightarrow \infty; n \neq m} \|x_n - x_m\|$ exists. Then*

$$\limsup_{n \rightarrow \infty} \|x_n\| \leq \lim_{n,m \rightarrow \infty; n \neq m} \|x_n - x_m\|.$$

Proposition 8 ([12]). *Let $\mathbf{a} = (a_j), \mathbf{b} = (b_j) \in \mathbb{R}^N$ and let a norm $\|\cdot\|_A$ on \mathbb{R}^N be monotone. If $\|\mathbf{a}\| = \|\mathbf{b}\|$, $|a_j| \leq |b_j|$ ($j = 1, \dots, N$) and $|a_{j_0}| < |b_{j_0}|$ then $\|(\chi_{N \setminus \{j_0\}}(j)a_j)\| = \|(\chi_{N \setminus \{j_0\}}(j)b_j)\|$, where $N = \{1, \dots, N\}$ and $\chi_{N \setminus \{j_0\}}$ is the characteristic function of $N \setminus \{j_0\}$.*

By Theorem 1, Propositions 5, 6, 7 and 8 we can prove the following key result.

Theorem 9. *Let X_1, \dots, X_N be Banach spaces and let a norm $\|\cdot\|_A$ on \mathbb{R}^N have Property T_1^N . Then $R(1, (X_1 \oplus \dots \oplus X_N)_A) < 2$ if and only if $R(1, X_j) < 2$ for all $1 \leq j \leq N$*

Corollary 10 (cf. [15]). *Let X and Y be Banach spaces and let a norm on \mathbb{R}^2 be not ℓ_1 -norm. Then $R(1, (X \oplus_A Y)) < 2$ if and only if $R(1, X) < 2$ and $R(1, Y) < 2$.*

Theorem 9 combined with Theorem 3 yields nontrivialness of Dominguez-Benavides coefficient of A -direct sums of N uniformly non-square Banach spaces.

Theorem 11. *Let X_1, \dots, X_N be uniformly non-square Banach spaces and let a norm $\|\cdot\|_A$ on \mathbb{R}^N have Property T_1^N . Then $R(1, (X_1 \oplus \dots \oplus X_N)_A) < 2$.*

By Theorem 11 and Theorem 2 we obtain our main results.

Theorem 12. *Let X_1, \dots, X_N be uniformly non-square Banach spaces and let a norm $\|\cdot\|_A$ on \mathbb{R}^N have Property T_1^N . Then $(X_1 \oplus \dots \oplus X_N)_A$ has the fixed point property for nonexpansive mappings.*

Theorem 13. Let X_1, \dots, X_N be uniformly non-square Banach spaces and let a norm $\|\cdot\|_A$ on \mathbb{R}^N have Property T_∞^N . Then $(X_1^* \oplus \dots \oplus X_N^*)_{A^*}$ has the fixed point property for nonexpansive mappings, where $(X_1^* \oplus \dots \oplus X_N^*)_{A^*}$ is an A -direct sum of X_1^*, \dots, X_N^* whose norm is defined by the dual norm $\|\cdot\|_{A^*}$.

Acknowledgments

The authors were supported in part by JSPS KAKENHI (C) Grant number JP26400131.

REFERENCES

- [1] J.M. Ayerbe, T. Domínguez Benavides and G. Lopez, *Measure of noncompactness in metric fixed point theory*, Birkhäuser, 1997.
- [2] R. Bhatia, *Matrix Analysis*, Springer-Verlag, New York, 1997.
- [3] Y. Cui, H. Hudzik, M. Wisła, *M-Constants, Dominguez-Benavides coefficient, and weak fixed point property in Orlicz sequence spaces equipped with the p -Amemiya norm*, Fixed Point Theory Appl. **2016:89**(2016), 1-14.
- [4] S. Dhompangsa, M. Kato and T. Tamura, Uniform non-squareness for A -direct sums of Banach spaces with a strictly monotone norm, Linear Nonlinear Analysis **1** (2015), 247-260.
- [5] T. Domínguez Benavides, *Some properties of the set and ball measures of noncompactness and applications*, J. London Math. Soc. **34** (1986), 120-128.
- [6] T. Domínguez Benavides, *A geometrical coefficient implying the fixed point property and stability results*, Houston J. Math. **22** (1996), 835-849.
- [7] P. N. Dowling and S. Saejung, Non-squareness and uniform non-squareness of Z -direct sums, J. Math. Anal. Appl. **369** (2010), 53-59.
- [8] J. Garcia-Falset, E. Llorens-Fuster and Eva M. Mazcunan-Navarro, Uniformly non-square Banach spaces have the fixed point property for nonexpansive mappings, J. Funct. Anal. **233** (2006), 494-514.
- [9] R. C. James, Uniformly non-square Banach spaces, Ann. of Math. **80** (1964), 542-550.
- [10] M. Kato, K.-S. Saito and T. Tamura, On the ψ -direct sums of Banach spaces and convexity, J. Aust. Math. Soc. **75** (2003), 413-422.
- [11] M. Kato, K.-S. Saito and T. Tamura, Uniform non- ℓ_1^n -ness of ψ -direct sums of Banach spaces, J. Nonlinear Convex Anal. **11** (2010), 13-33.
- [12] M. Kato and T. Tamura, Weak nearly uniform smoothness of the ψ -direct sums $(X_1 \oplus \dots \oplus X_N)_\psi$, Comment. Math. **52** (2012), 171-198.
- [13] T. R. Landes, Permanence properties of normal structure, Pacific J. Math. **10** (1984), 125-143.
- [14] K.-S. Saito, M. Kato and Y. Takahashi, On absolute norms on \mathbb{C}^n , J. Math. Anal. Appl. **252** (2000), 879-905.
- [15] T. Tamura, $M(X \oplus_\psi Y)$ for the ψ -direct sum of two Banach spaces X and Y , Studies on Humanities and Social Sciences of Chiba University **25**(2012), 1-9.
- [16] T. Tamura, On Dominguez-Benavides coefficient of ψ -direct sums $(X_1 \oplus \dots \oplus X_N)_\psi$ of Banach spaces, Linear and Nonlinear Analysis **3** (2017), 87-99.